

# Successively Thresholded Domain Boundary Roughening Driven by Pinning Centers and Missing Bonds: Hard-Spin Mean-Field Theory Applied to $d=3$ Ising Magnets

Tolga Çağlar<sup>1</sup> and A. Nihat Berker<sup>1,2</sup>

<sup>1</sup>*Faculty of Engineering and Natural Sciences, Sabancı University, Tuzla, Istanbul 34956, Turkey*

<sup>2</sup>*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

Hard-spin mean-field theory has recently been applied to Ising magnets, correctly yielding the absence and presence of an interface roughening transition respectively in  $d = 2$  and  $d = 3$  dimensions and producing the ordering-roughening phase diagram for isotropic and anisotropic systems. The approach has now been extended to the effects of quenched random pinning centers and missing bonds on the interface of isotropic and anisotropic Ising models in  $d = 3$ . We find that these frozen impurities cause domain boundary roughening that exhibits consecutive thresholding transitions as a function interaction of anisotropy. For both missing-bond and pinning-center impurities, for moderately large values the anisotropy, the systems saturate to the "solid-on-solid" limit, exhibiting a single universal curve for the domain boundary width as a function of impurity concentration.

PACS numbers: 68.35.Dv, 05.50.+q, 64.60.De, 75.60.Ch

## I. INTRODUCTION

Hard-spin mean-field theory [1, 2] has recently been applied to Ising magnets, correctly yielding the absence and presence of an interface roughening transition respectively in  $d = 2$  and  $d = 3$  dimensions and producing the ordering-roughening phase diagram for isotropic and anisotropic systems.[3] The approach is now extended to the effects of quenched random pinning centers and missing bonds on the interface of isotropic and uniaxially anisotropic Ising models in  $d = 3$ . We find that these frozen impurities cause domain boundary roughening that exhibits consecutive thresholding transitions as a function interaction of anisotropy. We also find that, for moderately large values the anisotropy, the systems saturate to the "solid-on-solid" limit, exhibiting a single universal curve for the domain boundary width as a function of impurity concentration.

## II. THE ANISOTROPIC $d = 3$ ISING MODEL WITH IMPURITIES AND HARD-SPIN MEAN-FIELD THEORY

### A. The $d=3$ Anisotropic Ising Model

The  $d = 3$  anisotropic Ising model is defined by the Hamiltonian

$$-\beta\mathcal{H} = J_{xy} \sum_{\langle ij \rangle}^{xy} s_i s_j + J_z \sum_{\langle ij \rangle}^z s_i s_j, \quad (1)$$

where at each site  $i$  of a cubic lattice, the spin takes on the values  $s_i = \pm 1$ . The first sum is over the nearest-neighbor pairs of sites along the  $x$  and  $y$  spatial directions and the second sum is over the nearest-neighbor pairs of sites along the  $z$  spatial direction. The system has ferromagnetic interactions  $J_{xy}, J_z > 0$ , periodic boundary conditions in the  $x$  and  $y$  directions and oppositely




FIG. 1: A  $yz$  plane at temperature  $1/J_{xy} = 0.1$ . Filled and empty circles respectively represent the calculated local magnetizations with  $m_i > 0$  and  $m_i < 0$ . The left side is for the pure system,  $p = 0$ . The right side is calculated with quenched random pinning centers with concentration  $p = 0.24$ . Islands that are disconnected from the pinned  $z$  boundary plane of their own sign (typically occurring around an opposite pinning center deep inside a bulk phase) do not enter the interface width calculation and are not shown here. Thus, the disconnected pieces seen in this figure are actually part of an overhang, connected to the corresponding  $z$  boundary plane via the other  $yz$  planes. The dashed lines delimit the domain boundary and the separation between these dashed lines gives the domain boundary width in this  $yz$  plane. The same procedure for determining the interface width is also applied to the missing bond systems.

fixed boundary conditions at the two terminal planes in the  $z$  spatial direction, which yields a domain boundary within the system when in the ordered phase. Thus, the system is generally uniaxially anisotropic. We systematically study the anisotropic  $J_{xy} \neq J_z$  as well as the isotropic  $J_{xy} = J_z$  cases.




FIG. 2: (Color online) Calculated domain boundary widths versus impurity concentration  $p$  for different anisotropy  $J_z/J_{xy}$  values, at temperature  $1/J_{xy} = 0.1$ . In the upper panel for missing bonds, from the bottom to the top curves, the anisotropies are  $J_z/J_{xy} = 0.1$  to  $5.0$  with  $0.1$  intervals and  $J_z/J_{xy} = 5.5$  to  $10$  with  $0.5$  intervals. The dashed curves are calculated with the predicted threshold anisotropy values of  $J_z/J_{xy} = 1, 2, 3, 4, 5$ . In the lower panel for pinning centers, the anisotropies are  $J_z/J_{xy} = 0.5$  to  $2.5$  with  $0.1$  intervals. The dashed curves are calculated with the predicted threshold anisotropy values of  $J_z/J_{xy} = 1, 2$ . Beyond  $J_z/J_{xy} \simeq 5$  and  $2.3$ , respectively for missing bonds and pinning centers, the system saturates to the  $J_z/J_{xy} \rightarrow \infty$  "solid-on-solid" limit, exhibiting a single universal curve for the domain boundary width as a function of impurity concentration, for all  $J_z/J_{xy} \gtrsim 5$  and  $J_z/J_{xy} \gtrsim 2.3$  respectively.

### B. Method: Hard-Spin Mean-Field Theory

In our current study, hard-spin mean-field theory [1, 2], which has been qualitatively and quantitatively successful in frustrated and unfrustrated, equilibrium and non-equilibrium magnetic ordering problems [3–17], including recently the interface roughening transition [3], is used to study the roughening of an interface by quenched random pinning center sites or missing bonds. The self-consistency equations of hard-spin mean-field theory [2] are

$$m_i = \sum_{\{s_j\}} \left[ \left( \prod_j \frac{1 + m_j s_j}{2} \right) \tanh \left( \sum_j J_{ij} s_j \right) \right], \quad (2)$$




FIG. 3: (Color online) Calculated domain boundary widths versus anisotropy  $J_z/J_{xy}$ , at temperature  $1/J_{xy} = 0.1$ . The consecutive curves, bottom to top, are for impurity concentration values of  $p = 0.04$  to  $0.72$  (top panel) and  $1$  (bottom panel) with  $0.04$  intervals. These values of  $p$  are noted next to the curves. The curves show the deviations from the isotropic case  $J_z/J_{xy} = 1$  (vertical dash-dotted line) in the directions of strongly coupled planes  $J_z/J_{xy} > 1$  or weakly coupled planes  $J_z/J_{xy} < 1$ . The predicted threshold values are well reproduced by the calculated widths. It is clearly seen to the right of this figure that beyond  $J_z/J_{xy} \simeq 5$  and  $2.3$ , respectively for missing bonds and pinning centers, the system saturates to the  $J_z/J_{xy} \rightarrow \infty$  "solid-on-solid" limit, exhibiting a single universal value for the domain boundary width as a function of impurity concentration, for all  $J_z/J_{xy} \gtrsim 5$  and  $J_z/J_{xy} \gtrsim 2.3$  respectively.

where  $m_i = \langle s_i \rangle$  is the local magnetization at site  $i$ , the sum  $\{s_j\}$  is over all possible values of the spins  $s_j$  at the nearest-neighbor sites  $j$  to site  $i$ , and  $m_j$  are the magnetizations at the nearest-neighbor sites. These coupled equations for all sites are solved by local numerical iteration, in a  $10 \times 10 \times 10$  system.

### III. DOMAIN BOUNDARY WIDTHS

#### A. Determination of the Domain Boundary Width

In our study, the domain boundary is roughened in two ways: (1) Magnetic impurities are included in the system by pinning randomly chosen sites to  $s_i = +1$  or to  $s_i = -1$ . The impurity concentration  $p$  in this case is the ratio of the number of pinned sites to the total number of sites. The numbers of  $+1$  and  $-1$  pinned sites are fixed to be equal, to give both domains an equal chance to advance over its counter. (2) Missing bonds are created by removing randomly chosen bonds. In this case, the concentration  $p$  is given by the ratio of the number of removed bonds to the total number of bonds.

The domain boundary width is calculated by first considering each  $yz$  plane. The boundary width in each  $yz$  plane is calculated by counting the number of sites, in the  $z$  direction, between the two furthest opposite magnetizations in the plane (Fig. 1). This number is averaged over all the  $yz$  planes. The result is then averaged over 100 independent realizations of the quenched randomness.

#### B. Impurity Effects on the Domain Boundary Width

Our calculated domain boundary widths, as a function of impurity (*i.e.*, missing-bond or pinned-site) concentration  $p$  at temperature  $1/J_{xy} = 0.1$ , are shown in Fig. 2. The different curves are for different interaction anisotropies  $J_z/J_{xy}$ . In the lower panel for pinning-center impurity, the domain boundary roughens with the introduction of infinitesimal impurity, for all anisotropies: The curves have finite slope at the pure system. In the upper panel for missing-bond impurity, the domain boundary roughens with the introduction of infinitesimal impurity for strongly coupled planes  $J_z/J_{xy} > 2.5$ , whereas for weakly coupled planes  $J_z/J_{xy} < 2.5$ , it is seen that infinitesimal or small impurity has essentially no effect on the flat domain boundary. In the latter cases, the curves reach the pure system with zero slope.

For both missing-bond and pinning-center impurities, for moderately large values of  $J_z/J_{xy}$ , we find (Figs. 2 and 3) that the systems saturate to the  $J_z/J_{xy} \rightarrow \infty$  "solid-on-solid" limit [18]. Thus, the systems exhibit a single universal curve for the domain boundary width as a function of impurity concentration, onwards from all moderately large values of  $J_z/J_{xy}$ .

#### C. Successive Roughening Thresholds

A bunching of the curves is visible, in the domain-boundary width curves in Fig. 2, especially in the upper panel for missing-bond impurity. This corresponds to a thresholded domain boundary roughening, controlled by the interaction anisotropy. This effect is also visible in

Fig. 3, by the steps in the curves which give the domain boundary widths as a function of the interaction anisotropy  $J_z/J_{xy}$  for different impurity concentrations  $p$ , at temperature  $1/J_{xy} = 0.1$ .

Thresholded domain boundary roughening can be understood by considering the effect of increasing the anisotropy. We first discuss the case of missing-bond impurity. Upon increasing  $J_z$ , for what value of  $J_z$  will a spin flip, *e.g.*, from  $+1$  to  $-1$ , thereby increasing the domain boundary width (directly and/or by inducing a flip cascade)? Increasing  $J_z$  can flip a spin and increase the width only if one of its bonds in the  $\pm z$  direction is missing and the non-missing bond connects to a  $-1$  spin. This flip will then happen for  $J_z = (q - q')J_{xy}$ , where  $(q, q')$  are the numbers of  $xy$  neighbors bonded to the flipping spin that are respectively  $+1, -1$ . The possible values are  $(q, q') = (4, 0), (3, 0), (2, 0), (1, 0), (3, 1), (2, 1)$ , giving the threshold values of  $J_z/J_{xy} = 1, 2, 3, 4$ , in fact calculationally seen in the top panels of Figs. 2 and 3. Furthermore, the simultaneous flip of two neighboring spins gives the threshold value of  $J_z/J_{xy} = 5$ , also calculationally seen in the top panels of Figs. 2 and 3. Beyond  $J_z/J_{xy} = 5$ , the system saturates to the  $J_z/J_{xy} \rightarrow \infty$  "solid-on-solid" limit [18], exhibiting a single universal curve for the domain boundary width as a function of impurity concentration, for all  $J_z/J_{xy} \gtrsim 5$ .

We now discuss the case of pinned-site impurity. We again consider the effect of increasing  $J_z$  and investigate the value of  $J_z$  that will flip the spin, *e.g.*, from  $+1$  to  $-1$ , thereby increasing the domain boundary width (again, directly and/or by inducing a flip cascade). Increasing  $J_z$  can flip this spin only if both of its neighbors in the  $\pm z$  direction are  $-1$ , with one of these being part of a disconnected island seeded by a pinning center. This flip will then happen for  $2J_z = (q - q')J_{xy}$ , where again  $q$  and  $q'$  are the numbers of  $xy$  neighbors bonded to the flipping spin that are respectively  $+1$  and  $-1$ . The possible values are  $(q, q') = (4, 0), (3, 1)$ , giving the threshold values of  $J_z/J_{xy} = 1, 2$ , calculationally seen in the bottom panels of Figs. 2 and 3. Beyond  $J_z/J_{xy} \simeq 2.3$ , the system saturates to the  $J_z/J_{xy} \rightarrow \infty$  "solid-on-solid" limit [18], exhibiting a single universal curve for the domain boundary width as a function of impurity concentration, for all  $J_z/J_{xy} \gtrsim 2.3$ .

On a similar vein, in the limit of  $xy$  planes weakly coupled due to low  $J_z/J_{xy}$  and high concentration of missing bonds, the domain boundary gains by the intermediacy of sending overhangs in the lateral  $x$  and  $y$  directions, eventually covering the whole system via randomly magnetized  $xy$  planes. In this case, the spin is flipped by the effect of  $J_{xy}$  upon decreasing  $J_z$ . This flip occurs at  $2J_z = (q - q')J_{xy}$ , where  $(q, q')$  has to be such that  $J_z/J_{xy}$  is low. Thus,  $(q, q') = (2, 1)$ . (Other pairs of values,  $(3, 0)$  and  $(1, 0)$  do not contribute to this spread of overhangs.) Indeed, in Fig. 3, a rise in the domain for decreasing  $J_z < 0.5$  is seen at high missing bond concentration.

#### IV. CONCLUSION

The effects of quenched random pinning centers and missing bonds on the interface of isotropic and uniaxially anisotropic Ising models in  $d = 3$  have been investigated by hard-spin mean-field theory. We find that the frozen impurities cause domain boundary roughening that exhibits consecutive thresholding transitions as a function of interaction of anisotropy  $J_z/J_{xy}$ . The numerical results, showing the thresholding transitions as the bunching of domain boundary width versus impurity concentration curves (Fig. 2) and steps in the domain boundary width versus anisotropy curves (Fig. 3) agree with our spin-flip arguments at the interface. For both missing-bond and

pinning-center impurities, for moderately large values of  $J_z/J_{xy}$ , the systems saturate to the  $J_z/J_{xy} \rightarrow \infty$  "solid-on-solid" limit, thus exhibiting a single universal curve for the domain boundary width as a function of impurity concentration, onwards from all moderately large values of  $J_z/J_{xy}$ .

#### Acknowledgments

Support by the Alexander von Humboldt Foundation, the Scientific and Technological Research Council of Turkey (TÜBİTAK), and the Academy of Sciences of Turkey (TÜBA) is gratefully acknowledged.

- 
- [1] R. R. Netz and A. N. Berker, Phys. Rev. Lett. **66**, 377 (1991).
  - [2] R. R. Netz and A. N. Berker, J. Appl. Phys. **70**, 6074 (1991).
  - [3] T. Çağlar and A. N. Berker, Phys. Rev. E **84**, 051129 (2011).
  - [4] J. R. Banavar, M. Cieplak, and A. Maritan, Phys. Rev. Lett. **67**, 1807 (1991).
  - [5] R. R. Netz and A. N. Berker, Phys. Rev. Lett. **67**, 1808 (1991).
  - [6] R. R. Netz, Phys. Rev. B **46**, 1209 (1992).
  - [7] R. R. Netz, Phys. Rev. B **48**, 16113 (1993).
  - [8] A. N. Berker, A. Kabakçioğlu, R. R. Netz, and M. C. Yalabık, Turk. J. Phys. **18**, 354 (1994).
  - [9] A. Kabakçioğlu, A. N. Berker, and M. C. Yalabık, Phys. Rev. E **49**, 2680 (1994).
  - [10] E. A. Ames and S. R. McKay, J. Appl. Phys. **76**, 6197 (1994).
  - [11] G. B. Akgüç and M. C. Yalabık, Phys. Rev. E **51**, 2636 (1995).
  - [12] J. E. Tesiero and S. R. McKay, J. Appl. Phys. **79**, 6146 (1996).
  - [13] J. L. Monroe, Phys. Lett. A **230**, 111 (1997).
  - [14] A. Pelizzola and M. Pretti, Phys. Rev. B **60**, 10134 (1999).
  - [15] A. Kabakçioğlu, Phys. Rev. E **61**, 3366 (2000).
  - [16] H. Kaya and A. N. Berker, Phys. Rev. E **62**, R1469 (2000); also see M. D. Robinson, D. P. Feldman, and S. R. McKay, Chaos **21**, 037114 (2011).
  - [17] O. S. Sariyer, A. Kabakçioğlu, and A. N. Berker, Phys. Rev. E **86**, 041107 (2012).
  - [18] R. H. Swendsen, Phys. Rev. B **15**, 689 (1977).